

Goat Anti-Rhesus IgG (H+L) Secondary Antibody

Goat Anti-Rhesus IgG (H+L)-HRP


CATALOG # AAA78849	HOST Goat	APPLICATIONS Western Blot, Immunohistochemistry, Immunofluorescence, Immunoblot, Western Blot, ELISA
------------------------------	---------------------	--

PRODUCT SYNONYMS

Goat Anti-Rhesus IgG (H+L); Polyclonal Secondary Antibody; Goat Anti-Rhesus IgG (H+L)-HRP; Goat Anti-Rhesus IgG(H+L) - HRP; Goat Anti-Rhesus IgG (H+L) secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Specificity	Reacts with the heavy and light chains of rhesus IgG.
Purity/Purification	Affinity chromatography on rhesus IgG covalently linked to agarose.
Form/Format	Horseradish Peroxidase (HRP) Conjugate
Applicable Applications for Goat Anti-Rhesus IgG (H+L) secondary antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), IB (Immunoblot), ELISA
Source	Pooled antisera from goats hyperimmunized with rhesus IgG.
Cross Absorption	None; may react with immunoglobulins from other species and the light chains of other rhesus immunoglobulins.
Characterization	To ensure lot-to-lot consistency, each batch of product is tested by ELISA and/or Fluorescent ELISA for conformance to characteristics of a standard reference reagent.
Preparation and Storage	The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of 100 mM borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added. Store at 2-8 degree C. The fluorescein (FITC) and rhodamine (TRITC) conjugates are supplied as 1.0 mg in 1.0 mL PBS/NaN₃. Store at 2-8 degree C. The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL of stock solution in 50mM Tris/1mM MgCl₂/50% Glycerol, pH 8.0, containing 0.1% NaN₃ as preservative. Store at 2-8 degree C or long-term at -20 degree C. The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8 degree C or long-term at -20 degree C. The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL PBS/NaN₃. Store at 2-8 degree C. Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.



Application Data
FLISA plate was coated with purified rhesus IgG. Immunoglobulin was detected with Goat Anti-Rhesus IgG (H+L)-PE

ADDITIONAL INFORMATION

Product Categories/Family for Goat Anti-Rhesus IgG (H+L) secondary antibody	Goat Anti-Rhesus IgG (H+L)
---	----------------------------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/goat-anti-rhesus-igg-hl-secondary-antibody-western-blot-immunohistochemistry-immunofluorescence-immunoblot-elisa/78849>