

Goat Anti-Human IgG Fc, Bov/Hrs/Mse SP. Ads Secondary Antibody

Goat Anti-Human IgG Fc, Bov/Hrs/Mse SP. Ads-UNLB

CATALOG # AAA78880	HOST Goat	REACTIVITY Human	APPLICATIONS ELISA
------------------------------	---------------------	----------------------------	------------------------------

PRODUCT SYNONYMS

Goat Anti-Human IgG Fc; Bov/Hrs/Mse SP. Ads; Polyclonal Secondary Antibody; Bov/Hrs/Mse SP. Ads-UNLB; Bov/Hrs/Mse SP. Ads secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	Reacts with the Fc region of human IgG
Purity/Purification	Affinity chromatography on human IgG Fc covalently linked to agarose
Form/Format	The purified (UNLB) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added.
Applicable Applications for Goat Anti-Human IgG Fc, Bov/Hrs/Mse SP. Ads secondary antibody	ELISA
Cross Absorption	Human IgG Fab, IgM, and IgA; mouse, bovine, and horse serum proteins (SP); may react with IgG from other species
Source Note	Pooled antisera from goats hyperimmunized with normal human IgG and human IgG paraproteins
Preparation and Storage	Store at 2-8°C. Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

ADDITIONAL INFORMATION

Product Categories/Family for Goat Anti-Human IgG Fc, Bov/Hrs/Mse SP. Ads secondary antibody	Secondary Antibodies & Reagents - Polyclonal
--	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/goat-anti-human-igg-fc-bov-hrs-mse-sp-ads-secondary-antibody-elisa/78880>