

Mouse anti-Human VEGF-A Monoclonal Antibody | anti-VEGF-A antibody

Mouse Anti-Human VEGF-A

CATALOG # AAA79102	HOST Mouse	REACTIVITY Human	APPLICATIONS Western Blot, ELISA
------------------------------	----------------------	----------------------------	--


PRODUCT SYNONYMS

VEGF-A; Monoclonal Antibody; Mouse Anti-Human VEGF-A; Anti-human VEGF-A (#339/H2); Vascular endothelial growth factor-A; VPF; anti-VEGF-A antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Clone Number	IgG1; 339/H2
Purity/Purification	Protein G purified
Form/Format	Lyophilized
Sequence Length	412
Applicable Applications for anti-VEGF-A antibody	WB (Western Blot), ELISA
Immunogen	recombinant human VEGF189 (RT# 300-094)
Preparation	Monoclonals were produced with the help of BALB/c mice using recombinant human VEGF189 derived from E. coli.
Buffer	PBS
Preparation and Storage	The lyophilized antibody is stable at room temperature for up to 1 month. The reconstituted antibody is stable for at least two weeks at 2-8 degree C. Frozen aliquots are stable for at least 6 months when stored at -20 degree C.

IMAGES



Application Data



Application Data

ADDITIONAL INFORMATION

Related Product Information for anti-VEGF-A antibody	Vascular endothelial growth factor (VEGF or VEGF-A), also known as vascular permeability factor (VPF) or vasculotropin, is a homodimeric 34 - 42 kDa, heparin-binding glycoprotein with potent angiogenic, mitogenic and vascular permeability-enhancing activities specific for endothelial cells. Different isoforms can be generated by differential splicing (e.g. VEGF165). All eight cysteine residues involved in intra- and inter-chain disulfide bonds are conserved among these growth factors. A cDNA encoding a protein having a 53% amino acid sequence homology in the PDGF-like region of VEGF has been isolated from a human placental cDNA library. This protein, named placenta growth factor (PlGF), is now recognized to be a member of the VEGF family of growth factors. Two receptor tyrosine kinases have been described as putative VEGF receptors. Flt-1 (fms-like tyrosine kinase), and KDR (kinase-insert-domain-containing receptor) proteins have been shown to bind VEGF-A with high affinity. In vitro, VEGF is a potent endothelial cell mitogen. In cultured endothelial cells, VEGF can activate phospholipase C and induce rapid increases of free cytosolic Ca ²⁺ . VEGF has also been shown to be chemotactic for monocytes and osteoblasts. In vivo, VEGF can induce angiogenesis as well as increase microvascular permeability. As a vascular permeability factor, VEGF acts directly on the endothelium and does not degranulate mast cells. Based on its in vitro and in vivo properties, VEGF is expected to play important roles in inflammation and during normal and pathological angiogenesis, a process that is associated with wound healing, embryonic development, and growth and metastasis of solid tumors.
References	1. Breier et al., Dev 114:521, 1992 2. Fiebig et al., Eur J Biochem 211:19, 1993 3. Flamme et al., Dev Biol 162:699, 1995 4. Kremer et al., Cancer Res 57:3852, 1997

NCBI AND UNIPROT INFORMATION

NCBI GI #	76781480
NCBI GeneID	7422
NCBI Accession #	NP_001165095
NCBI GenBank Nucleotide #	NM_001171624
UniProt Accession #	P15692; O60720; O75875; Q074Z4; Q16889; Q5UB46; Q6P0P5; Q96KJ0; Q96L82; Q96NW5; Q9H1W8; B5BU86
Molecular Weight	27,042 Da
NCBI Official Full Name	vascular endothelial growth factor A isoform a
NCBI Official Synonym Full Names	vascular endothelial growth factor A
NCBI Official Symbol	VEGFA
NCBI Official Synonym Symbols	VPF; VEGF; MVCD1
NCBI Protein Information	vascular endothelial growth factor A; vascular permeability factor
UniProt Protein Name	Vascular endothelial growth factor A
UniProt Gene Name	VEGFA
UniProt Synonym Gene Names	VEGF; VEGF-A; VPF
UniProt Entry Name	VEGFA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/vegf-a-antibody-clone-igg1-339-h2-western-blot-elisa/79102>