

VEGF165 active protein

Human VEGF165

CATALOG # AAA79288	HOST E Coli	REACTIVITY Human
------------------------------	-----------------------	----------------------------


PRODUCT SYNONYMS

VEGF165; Human VEGF165; VEGF-A; VPF; VEGF165 active protein

PRODUCT SPECIFICATIONS

Host	E Coli
Reactivity	Human
Purity/Purification	> 95% by SDS-PAGE & silver stain
Form/Format	Sterile filtered protein solution was lyophilized from 50 mM acetic acid, free from stabilizers and preservatives.
Sequence Positions	APMAEGGGQNHHEVVKFMDVYQRSYCHPIETLVDIFQEYPDEIEYIFKPSCVPLMRCGGCCNDEGLECVPTESNITMQIMRIKPHQGQHIGEMSFLQHN
Sequence	N-Terminal Sequence:A PMAEGG
Sequence Length	165
Amino Acids	165 aa
Reconstitution	The lyophilized VEGF165 should be reconstituted in 50 mM acetic acid to a concentration not lower than 50 ug/ml. Store at 2°C to 8°C for up to 1 week
Buffer	50 mM acetic acid
Preparation and Storage	After initial reconstitution, the addition of at least 0.1% human or bovine serum albumin is recommended for frozen aliquots. Prepare and store working Product Form Temperature Storage Time Lyophilized -20°C to -80°C 1 year Lyophilized 2°C to 8°C 6 months Lyophilized Room Temperature 1 month

IMAGES



Application Data

Fig. 3: Confluent PAE/KDR cells were stimulated with 50ng/ml human VEGF165 for 10 min at 37°C. Cells were lysed and an IP was performed using a mouse anti-human KDR antibody (Cat# WB was performed with a mouse anti-human KDR antibody (Cat# and an anti-Phosphotyrosine antibody.





Figure 2. VEGF₁₆₅-induced proliferation of HUVE cells. HUVECs were stimulated with increasing amounts of human VEGF₁₆₅.

Application Data



Application Data

ADDITIONAL INFORMATION

Related Product Information for VEGF165 active protein	Human Vascular Endothelial Growth Factor VEGF165, a 23 kDa protein consisting of 165 amino acid residues, is produced as a homodimer. VEGF is a polypeptide growth factor and a member of the platelet-derived growth factor family. It is a specific mitogen for vascular endothelial cells and a strong angiogenic factor in vivo. Two high-affinity tyrosine kinase receptors for VEGF165 have been identified, VEGFR-1 (FLT-1), and VEGFR-2 (KDR). Consistent with the endothelial cell-specific action of VEGF165, expression of both receptor genes has been found predominantly but not exclusively on endothelial cells. Expression of VEGFR-1 was also found on human monocytes, neutrophils (PMNs), bovine brain pericytes and villous and extra villous trophoblast. In addition to its action as a mitogen it is a potent vascular permeability factor (VPF) in vivo. VEGF165 is also a chemo attractant molecule for monocytes and endothelial cells. 5 different proteins are generated by differential splicing: VEGF121, VEGF145, VEGF165, VEGF189 and VEGF206. The most abundant form is VEGF165. Whereas VEGF121 and VEGF165 are secreted proteins, VEGF145, VEGF189 and VEGF206 are strongly cell-associated. The isoforms VEGF145, VEGF165 and VEGF189 bind to heparin with high affinity. VEGF165 is apparently a homo-dimer, but preparations of VEGF165 show some heterogeneity on SDS gels, depending on the secretion of different glycosylation patterns. All dimeric forms have similar biological activities but their bioavailability is very different. There is good evidence that different cells and tissues express different VEGF isoforms. The other members of this increasing growth factor family are VEGF-B, -C, -D and -E. Another member is the Placenta growth factor PlGF.
Product Categories/Family for VEGF165 active protein	Cytokines & Growth Factors
References	1. Breier et al., Dev 114:521, 1992 2. Fiebig et al., Eur J Biochem 211:19, 1993 3. Flamme et al., Dev Biol 162:699, 1995 4. Kremer at al., Cancer Res 57:3852, 1997

NCBI AND UNIPROT INFORMATION

NCBI GI #	76781480
NCBI GeneID	7422
NCBI Accession #	NP_001165097
NCBI GenBank Nucleotide #	NM_001171626
UniProt Accession #	P15692; O60720; O75875; Q074Z4; Q16889; Q5UB46; Q6P0P5; Q96KJ0; Q96L82; Q96NW5; Q9H1W8; B5BU86
Molecular Weight	38,2 kDa
NCBI Official Full Name	vascular endothelial growth factor A isoform a
NCBI Official Synonym Full Names	vascular endothelial growth factor A
NCBI Official Symbol	VEGFA
NCBI Official Synonym Symbols	VPF; VEGF; MVCD1
NCBI Protein Information	vascular endothelial growth factor A; vascular permeability factor
UniProt Protein Name	Vascular endothelial growth factor A

UniProt Gene Name	VEGFA
UniProt Synonym Gene Names	VEGF; VEGF-A; VPF
UniProt Entry Name	VEGFA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/e-coli-human-active-protein/veg165/79288>