

Mouse cytochrome c, somatic ELISA Kit | CYCS elisa kit

Mouse Cytochrome-C, Cyt-C ELISA Kit

CATALOG #
AAA79883

REACTIVITY
Mouse


PRODUCT SYNONYMS

cytochrome c; somatic; Mouse Cytochrome-C; Cyt-C ELISA Kit; CYC; HCS; THC4; OTTHUMP00000202043; OTTHUMP00000202044; CYCS elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of mouse Cyt-C. No significant cross-reactivity or interference between mouse Cyt-C and analogues was observed.
Sequence Length	105
Samples	Serum, plasma, tissue homogenates, Cell lysates
Assay Type	Sandwich
Detection Range	34.3 pg/ml -2200 pg/ml.
Sensitivity	The minimum detectable dose of mouse Cyt-C is typically less than 8.5 pg/ml. The sensitivity of this assay, or Lower Limit of Detection (LLD) was defined as the lowest protein concentration that could be differentiated from zero. It was determined the mean O.D value of 20 replicates of the zero standard added by their three standard deviations.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): CV%<8%. Three samples of known concentration were tested twenty times on one plate to assess.
Inter-assay Precision	Inter-assay Precision (Precision between assays): CV%<10%. Three samples of known concentration were tested in twenty assays to assess.
Preparation and Storage	Unopened test kits should be stored at 2 to 8 degree C upon receipt. Please refer to pdf manual for further storage instructions.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CYCS elisa kit	Principle of the assay: This assay employs the quantitative sandwich enzyme immunoassay technique. Antibody specific for Cyt-C has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any Cyt-C present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for Cyt-C is added to the wells. After washing, avidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of Cyt-C bound in the initial step. The color development is stopped and the intensity of the color is measured.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	119388072
UniProt Accession #	P00008
Molecular Weight	11,692 Da
NCBI Official Full Name	Cytochrome c
UniProt Protein Name	Cytochrome c
UniProt Gene Name	CYCS
UniProt Synonym Gene Names	CYC
UniProt Entry Name	CYC_RABIT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/cycs-mouse-elisa-kits/cytochrome-c-somatic/79883>