

Human CD40 ligand ELISA Kit | CD40LG elisa kit

Human Soluble Cluster of differentiation 40 ligand, sCD40L ELISA Kit

CATALOG #
AAA80061

REACTIVITY
Human


PRODUCT SYNONYMS

CD40 ligand; Human Soluble Cluster of differentiation 40 ligand; sCD40L ELISA Kit; RP3-527F8.3; CD154; CD40L; HIGM1; IGM; IMD3; T-BAM; TNFSF5; TRAP; gp39; hCD40L; CD40 antigen ligand; T-B cell-activating molecule; TNF-related activation protein; tumor necrosis factor (ligand) superfa; CD40LG elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of human sCD40L. No significant cross-reactivity or interference between human sCD40L and analogues was observed.
Sequence Length	261
Assay Type	Quantitative Sandwich
Samples	Serum, plasma, tissue homogenates
Detection Range	0.312 ng/ml-20 ng/ml
Sensitivity	< 0.078 ng/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): CV%<8%. Three samples of known concentration were tested twenty times on one plate to assess.
Inter-assay Precision	Inter-assay Precision (Precision between assays): CV%<10%. Three samples of known concentration were tested in twenty assays to assess.
Preparation and Storage	Unopened test kits should be stored at 2 to 8 degree C upon receipt. Please refer to pdf manual for further storage instructions.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CD40LG elisa kit	Principle of the Assay: This assay employs the quantitative sandwich enzyme immunoassay technique. Antibody specific for sCD40L has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any sCD40L present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for sCD40L is added to the wells. After washing, avidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of sCD40L bound in the initial step. The color development is stopped and the intensity of the color is measured.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557433
NCBI GeneID	959
NCBI Accession #	NP_000065.1
NCBI GenBank Nucleotide #	NM_000074.2
UniProt Accession #	P29965
Molecular Weight	29,274 Da
NCBI Official Full Name	CD40 ligand
NCBI Official Synonym Full Names	CD40 ligand
NCBI Official Symbol	CD40LG
NCBI Official Synonym Symbols	IGM; IMD3; TRAP; gp39; CD154; CD40L; HIGM1; T-BAM; TNFSF5; hCD40L
NCBI Protein Information	CD40 ligand; CD40-L; CD40 antigen ligand; T-cell antigen Gp39; T-B cell-activating molecule; TNF-related activation protein; tumor necrosis factor (ligand) superfamily member 5
UniProt Protein Name	CD40 ligand
UniProt Gene Name	CD40LG
UniProt Synonym Gene Names	CD40L; TNFSF5; TRAP; CD40-L; TRAP
UniProt Entry Name	CD40L_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/cd40lg-human-elisa-kits/cd40-ligand/80061>