

Mouse chemokine (C-X3-C motif) ligand 1 ELISA Kit | CX3CL1 elisa kit

Mouse Fractalkine, FK ELISA Kit

CATALOG #
AAA80303

REACTIVITY
Mouse


PRODUCT SYNONYMS

chemokine (C-X3-C motif) ligand 1; Mouse Fractalkine; FK ELISA Kit; ABCD-3; C3Xkine; CXC3; CXC3C; NTN; NTT; SCYD1; fractalkine; neurotactin; small inducible cytokine subfamily D (Cys-X3-Cys); member 1 (fractalkine; neurotactin) ; small inducible cytokine subfamily D (; CX3CL1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Mouse CX3CL1. No significant cross-reactivity or interference between Mouse CX3CL1 and analogues was observed.
Sequence Length	395
Samples	Serum, plasma, cell culture supernates, tissue homogenates
Detection Range	125 pg/ml -8000 pg/ml
Sensitivity	31.25 pg/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): CV% is less than 8% Three samples of known concentration were tested twenty times on one plate to assess. Inter-assay Precision (Precision between assays): CV% is less than 10% Three samples of known concentration were tested in twenty assays to assess.
Detection Wavelength	450 nm
Sample Volume	50-100ul
Protein Biological Process 1	Cell Adhesion
Protein Biological Process 3	Cell adhesion
Preparation and Storage	Unopened test kits should be stored at 2 to 8 degree C upon receipt. Please refer to pdf manual for further storage instructions.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CX3CL1 elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to Fkn. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific for Fkn and Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then a TMB (3,3',5,5' tetramethyl-benzidine) substrate solution is added to each well. Only those wells that contain Fkn, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of a sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450 nm +/- 2 nm. The concentration of Fkn in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	31982017
NCBI GeneID	20312
NCBI Accession #	NP_033168.2
NCBI GenBank Nucleotide #	NM_009142.3
UniProt Accession #	O35188; O35933; Q91V44
Molecular Weight	42,099 Da
NCBI Official Full Name	fractalkine
NCBI Official Synonym Full Names	chemokine (C-X3-C motif) ligand 1
NCBI Official Symbol	Cx3cl1
NCBI Official Synonym Symbols	CX3C; Cxc3; Scyd1; ABCD-3; AB030188; AI848747; D8Bwg0439e
NCBI Protein Information	fractalkine; neurotactin; C-X3-C motif chemokine 1; small-inducible cytokine D1; CX3C membrane-anchored chemokine; small inducible cytokine subfamily D, 1
UniProt Protein Name	Fractalkine
UniProt Gene Name	Cx3cl1
UniProt Synonym Gene Names	Cx3c; Fkn; Scyd1
UniProt Entry Name	X3CL1_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/cx3cl1-mouse-elisa-kits/chemokine-c-x3-c-motif-ligand-1/80303>