

Goat interleukin 2 ELISA Kit | IL2 elisa kit

Goat Interleukin 2, IL-2 ELISA Kit

CATALOG #
AAA80366

REACTIVITY
Goat


PRODUCT SYNONYMS

interleukin 2; Goat Interleukin 2; IL-2 ELISA Kit; Goat Interleukin 2 (IL-2) ELISA Kit; IL-2; TCGF; lymphokine; T cell growth factor; aldesleukin; interleukin-2; involved in regulation of T-cell clonal expansion; IL2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Goat
Specificity	This assay has high sensitivity and excellent specificity for detection of goat IL-2. No significant cross-reactivity or interference between goat IL-2 and analogues was observed.
Sequence Length	155
Samples	Serum, plasma, tissue homogenates
Assay Type	Sandwich
Detection Range	15.6 pg/ml -1000 pg/ml.
Sensitivity	The minimum detectable dose of goat IL-2 is typically less than 3.9 pg/ml. The sensitivity of this assay, or Lower Limit of Detection (LLD) was defined as the lowest protein concentration that could be differentiated from zero. It was determined the mean O.D value of 20 replicates of the zero standard added by their three standard deviations.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): CV%<8%. Three samples of known concentration were tested twenty times on one plate to assess.
Inter-assay Precision	Inter-assay Precision (Precision between assays): CV%<10%. Three samples of known concentration were tested in twenty assays to assess.
Preparation and Storage	Unopened test kits should be stored at 2 to 8 degree C upon receipt. Please refer to pdf manual for further storage instructions.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL2 elisa kit	Principle of the assay: This assay employs the quantitative sandwich enzyme immunoassay technique. Antibody specific for IL-2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any IL-2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for IL-2 is added to the wells. After washing, avidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of IL-2 bound in the initial step. The color development is stopped and the intensity of the color is measured.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	547715
UniProt Accession #	P36835; P79156
Molecular Weight	17,704 Da
NCBI Official Full Name	Interleukin-2
UniProt Protein Name	Interleukin-2
UniProt Gene Name	IL2
UniProt Synonym Gene Names	IL-2; TCGF
UniProt Entry Name	IL2_CAPHI

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/il2-goat-elisa-kits/interleukin-2/80366>