

Human hepatocyte growth factor (hepapoietin A; scatter factor) ELISA Kit | HGF elisa kit

Human hepatocyte growth factor, HGF ELISA Kit

CATALOG #
AAA80738

REACTIVITY
Human


PRODUCT SYNONYMS

hepatocyte growth factor (hepapoietin A; scatter factor); Human hepatocyte growth factor; HGF ELISA Kit; HGF ELISA kit; DFNB39; F-TCF; HGFB; HPTA; SF; OTTHUMP00000206710; fibroblast-derived tumor cytotoxic factor; hepatocyte growth factor; lung fibroblast-derived mitogen; HGF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Human HGF. No significant cross-reactivity or interference between Human HGF and analogues was observed.
Sequence Length	728
Samples	Serum, plasma, tissue homogenates
Detection Range	6.25 pg/ml-400 pg/ml
Sensitivity	1.56 pg/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): CV% is less than 8% Three samples of known concentration were tested twenty times on one plate to assess. Inter-assay Precision (Precision between assays): CV% is less than 10% Three samples of known concentration were tested in twenty assays to assess.
Detection Wavelength	450 nm
Sample Volume	50-100ul
Preparation and Storage	Unopened test kits should be stored at 2 to 8 degree C upon receipt. Please refer to pdf manual for further storage instructions.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HGF elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to HGF. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific for HGF and Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then a TMB (3,3',5,5' tetramethyl-benzidine) substrate solution is added to each well. Only those wells that contain HGF, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of a sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450 nm +/- 2 nm. The concentration of HGF in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	33859835
NCBI GeneID	3082
NCBI Accession #	NP_000592.3
NCBI GenBank Nucleotide #	NM_000601.4
UniProt Accession #	P14210; Q02935; Q13494; Q14519; Q3KRB2; Q8TCE2; Q9BYL9; Q9BYM0; Q9UDU6; A1L3U6
Molecular Weight	83,134 Da
NCBI Official Full Name	hepatocyte growth factor isoform 1 preproprotein
NCBI Official Synonym Full Names	hepatocyte growth factor (hepapoietin A; scatter factor)
NCBI Official Symbol	HGF
NCBI Official Synonym Symbols	SF; HGFB; HPTA; F-TCF; DFNB39
NCBI Protein Information	hepatocyte growth factor; hepatopoeitin-A; hepatopoietin-A; lung fibroblast-derived mitogen; fibroblast-derived tumor cytotoxic factor
UniProt Protein Name	Hepatocyte growth factor
UniProt Gene Name	HGF
UniProt Synonym Gene Names	HPTA; SF
UniProt Entry Name	HGF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.