

Human chemokine (C-C motif) ligand 5 ELISA Kit | CCL5 elisa kit

Human C-C motif chemokine 5, CCL5/D17S136E/SCYA5 ELISA Kit

CATALOG #
AAA80893

REACTIVITY
Human


PRODUCT SYNONYMS

chemokine (C-C motif) ligand 5; Human C-C motif chemokine 5; CCL5/D17S136E/SCYA5 ELISA Kit; Human C-C motif chemokine 5 (CCL5/D17S136E/SCYA5) ELISA kit; D17S136E; MGC17164; RANTES; SCYA5; SISd; TCP228; OTTHUMP00000197106; SIS-delta; T-cell specific protein p288; beta-chemokine RANTES; regulated upon activation; normally T-expressed; and presumably secre; Human C-C motif chemokine 5 (CCL5/D17S136E/SCYA5); CCL5 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of human CCL5. No significant cross-reactivity or interference between human CCL5 and analogues was observed.
Sequence Length	91
Samples	Serum, plasma, cell culture supernates, tissue homogenates
Assay Type	Sandwich
Detection Range	15.6 pg/ml -1000 pg/ml.
Sensitivity	The minimum detectable dose of human CCL5 is typically less than 3.9 pg/ml. The sensitivity of this assay, or Lower Limit of Detection (LLD) was defined as the lowest protein concentration that could be differentiated from zero. It was determined the mean O.D value of 20 replicates of the zero standard added by their three standard deviations.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): CV%<8%. Three samples of known concentration were tested twenty times on one plate to assess.
Inter-assay Precision	Inter-assay Precision (Precision between assays): CV%<10%. Three samples of known concentration were tested in twenty assays to assess.
Preparation and Storage	Unopened test kits should be stored at 2 to 8 degree C upon receipt. Please refer to pdf manual for further storage instructions.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CCL5 elisa kit

Principle of the assay: This assay employs the quantitative sandwich enzyme immunoassay technique. Antibody specific for CCL5 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any CCL5 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for CCL5 is added to the wells. After washing, avidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of CCL5 bound in the initial step. The color development is stopped and the intensity of the color is measured.

NCBI AND UNIPROT INFORMATION

NCBI GI #	22538814
NCBI GeneID	6352
NCBI Accession #	NP_002976.2
NCBI GenBank Nucleotide #	NM_002985.2
UniProt Accession #	P13501; O43646; Q0QVW8; Q4ZGJ1; Q9NYA2; Q9UBG2; Q9UC99
Molecular Weight	9,990 Da
NCBI Official Full Name	C-C motif chemokine 5 isoform 1
NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 5
NCBI Official Symbol	CCL5
NCBI Official Synonym Symbols	SISd; eoCP; SCYA5; RANTES; TCP228; D17S136E; SIS-delta
NCBI Protein Information	C-C motif chemokine 5; beta-chemokine RANTES; T-cell specific protein p288; eosinophil chemotactic cytokine; small inducible cytokine subfamily A (Cys-Cys), member 5; regulated upon activation, normally T-expressed, and presumably secreted
UniProt Protein Name	C-C motif chemokine 5
UniProt Gene Name	CCL5
UniProt Synonym Gene Names	D17S136E; SCYA5; TCP228
UniProt Entry Name	CCL5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/ccl5-human-elisa-kits/chemokine-c-c-motif-ligand-5/80893>