

Bovine interleukin 22 ELISA Kit | IL22 elisa kit

Bovine interleukin 22, IL22 ELISA Kit

CATALOG #
AAA80948

REACTIVITY
Bovine


PRODUCT SYNONYMS

interleukin 22; Bovine interleukin 22; IL22 ELISA Kit; Bovine interleukin 22 (IL22) ELISA kit; IL-21; IL-22; IL-D110; IL-TIF; IL21; ILTIF; MGC79382; MGC79384; TIFIL-23; TIFa; zcyto18; IL-10-related T-cell-derived inducible factor; interleukin 21; IL22 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Bovine
Specificity	This assay has high sensitivity and excellent specificity for detection of Bovine IL22. No significant cross-reactivity or interference between Bovine IL22 and analogues was observed.
Sequence Length	179
Samples	Serum, plasma, tissue homogenates
Assay Type	Sandwich
Detection Range	15.6 pg/ml-1000 pg/ml
Sensitivity	3.9 pg/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): CV% is less than 8% Three samples of known concentration were tested twenty times on one plate to assess. Inter-assay Precision (Precision between assays): CV% is less than 10% Three samples of known concentration were tested in twenty assays to assess.
Detection Wavelength	450 nm
Sample Volume	50-100ul
Preparation and Storage	Unopened test kits should be stored at 2 to 8 degree C upon receipt. Please refer to pdf manual for further storage instructions.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL22 elisa kit	For SampleFor the quantitative determination of bovine interleukin 22 (IL22) concentrations in serum, plasma, tissue homogenates. Principle of the Assay: This assay employs the quantitative sandwich enzyme immunoassay technique. Antibody specific for IL22 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any IL22 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for IL22 is added to the wells. After washing, avidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of IL22 bound in the initial step. The color development is stopped and the intensity of the color is measured.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	10092625
NCBI GeneID	50616
NCBI Accession #	NP_065386.1
NCBI GenBank Nucleotide #	NM_020525.4
UniProt Accession #	Q9GZX6
Molecular Weight	20,011 Da
NCBI Official Full Name	interleukin-22
NCBI Official Synonym Full Names	interleukin 22
NCBI Official Symbol	IL22
NCBI Official Synonym Symbols	TIFa; IL-21; IL-22; ILTIF; IL-TIF; IL-D110; zcyto18; TIFIL-23
NCBI Protein Information	interleukin-22; cytokine Zcyto18; IL-10-related T-cell-derived inducible factor; IL-10-related T-cell-derived-inducible factor
UniProt Protein Name	Interleukin-22
UniProt Gene Name	IL22
UniProt Synonym Gene Names	ILTIF; ZCYTO18; UNQ3099/PRO10096; IL-22; IL-TIF
UniProt Entry Name	IL22_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.