

IgG Fab fragment Native Protein

Human IgG Fab fragment

CATALOG # AAA81642	HOST Purified from IgG	APPLICATIONS Western Blot, ELISA
------------------------------	----------------------------------	--


PRODUCT SYNONYMS

IgG Fab fragment; Human IgG Fab fragment; fragment antigen-binding; IgG Fab fragment native protein

PRODUCT SPECIFICATIONS

Host	Purified from IgG
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Liquid
Applicable Applications for IgG Fab fragment native protein	WB (Western Blot), ELISA
Peptide	Full length native Human IgG Fab fragment
Storage Buffer	pH7.4 PBS,50%glycerol
Preparation and Storage	Aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze / thaw cycles.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for IgG Fab fragment native protein	The fragment antigen-binding (Fab fragment) is a region on an antibody that binds to antigens. It is composed of one constant and one variable domain of each of the heavy and the light chain. These domains shape the paratope — the antigen-binding site — at the amino terminal end of the monomer. The two variable domains bind the epitope on their specific antigens.
---	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.