

UV excision repair protein RAD23 homolog A Recombinant Protein | RAD23A recombinant protein

Recombinant Human UV excision repair protein RAD23 homolog A

CATALOG # AAA81657	HOST E Coli or Yeast or Baculovirus or Mammalian Cell
------------------------------	---


PRODUCT SYNONYMS

UV excision repair protein RAD23 homolog A; Recombinant Human UV excision repair protein RAD23 homolog A; RAD23A recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli or Yeast or Baculovirus or Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	1-363aa; Full Length
Sequence	MAVTITLKTLLQQQTFKIRMEPDETVKVLKEKIEAEKGRDAFPVAGQKLIYAGKILSDDVPIRDYRIDEKNFVVVMVTKTKAGQGTSAPPEASPTAAPESSTSFPPAPTSGMSHPPPAAREDKSPSEESAPTTSPESVSGSVPSGSSGREEDAASTLVGTSEYETMLTEIMSMGYERERVVAALRASNNPHRAVEYLLTGIPGSPEPEHGSVQESQVSEQPATEAAGENPLEFLRDQPQFQNMQRVIQQLNPALLPALLQLGQENPQLLQQISRHQEQFIQMLNEPPGE LADISDVEGEVGAIGEEAPQMNYIQVTPQEKEAIERLKALGFPESLVIQAYFACEKNENLAANFLLSQNF DDE
Sequence Length	363
Production Note	Special Offer: The E Coli host-expressed protein is manufactured from a stock plasmid containing the protein gene. E Colihost-expressed protein is stocked in different unit sizes ranging from as small as 10 ug to as large as 1 mg. Bulk inventory is also available. The E Coli host-expressed protein has been ordered over and over again by researchers and has stood the test of time as both a robust protein and important target for the research community. It is part of our new program to make our most popular protein targets and corresponding hosts available in expanded unit sizes and with a quick processing time. Select E Coli host-expressed protein for the fastest delivery among all hosts. Please contact our or email to for more details.
Preparation and Storage	Store at -20 degree C, for extended storage, conserve at -20 degree C or -80 degree C.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for RAD23A recombinant protein	Multiubiquitin chain receptor involved in modulation of proteasomal degradation. Binds to 'Lys-48'-linked polyubiquitin chains in a length-dependent manner and with a lower affinity to 'Lys-63'-linked polyubiquitin chains. Proposed to be capable to bind simultaneously to the 26S proteasome and to polyubiquitinated substrates and to deliver ubiquitinated proteins to the proteasome. Involved in nucleotide excision repair and is thought to be functional equivalent for RAD23B in global genome nucleotide excision repair (GG-NER) by association with XPC. In vitro, the XPC:RAD23A dimer has NER activity. Can stabilize XPC. Involved in vpr-dependent replication of HIV-1 in non-proliferating cells and primary macrophages. Required for the association of HIV-1 vpr with the host proteasome.
Product Categories/Family for RAD23A recombinant protein	Epigenetics and Nuclear Signaling

References

- Purification and cloning of a nucleotide excision repair complex involving the Xeroderma pigmentosum group C protein and a human homologue of yeast RAD23. Masutani C., Sugasawa K., Yanagisawa J., Sonoyama T., Ui M., Enomoto T., Takio K., Tanaka K., van der Spek P.J., Bootsma D., Hoeijmakers J.H.J., Hanaoka F. *EMBO J.* 13:1831-1843(1994) NIEHS SNPs program Full-length cDNA libraries and normalization. Li W.B., Gruber C., Jessee J., Polayes D. Homo sapiens protein coding cDNA. Totoki Y., Toyoda A., Takeda T., Sakaki Y., Tanaka A., Yokoyama S., Ohara O., Nagase T., Kikuno R.F. The DNA sequence and biology of human chromosome 19. Grimwood J., Gordon L.A., Olsen A.S., Terry A., Schmutz J., Lamerdin J.E., Hellsten U., Goodstein D., Couronne O., Tran-Gyamfi M., Aerts A., Altherr M., Ashworth L., Bajorek E., Black S., Branscomb E., Caenepeel S., Carrano A.V., Caoile C., Chan Y.M., Christensen M., Cleland C.A., Copeland A., Dalin E., Dehal P., Denys M., Detter J.C., Escobar J., Flowers D., Fotopulos D., Garcia C., Georgescu A.M., Glavina T., Gomez M., Gonzales E., Groza M., Hammon N., Hawkins T., Haydu L., Ho I., Huang W., Israni S., Jett J., Kadner K., Kimball H., Kobayashi A., Larionov V., Leem S.-H., Lopez F., Lou Y., Lowry S., Malfatti S., Martinez D., McCready P.M., Medina C., Morgan J., Nelson K., Nolan M., Ovcharenko I., Pitluck S., Pollard M., Popkie A.P., Predki P., Quan G., Ramirez L., Rash S., Retterer J., Rodriguez A., Rogers S., Salamov A., Salazar A., She X., Smith D., Slezak T., Solovyev V., Thayer N., Tice H., Tsai M., Ustaszewska A., Vo N., Wagner M., Wheeler J., Wu K., Xie G., Yang J., Dubchak I., Furey T.S., DeJong P., Dickson M., Gordon D., Eichler E.E., Pennacchio L.A., Richardson P., Stubbs L., Rokhsar D.S., Myers R.M., Rubin E.M., Lucas S.M. *Nature* 428:529-535(2004) Human immunodeficiency virus type 1 Vpr interacts with HHR23A, a cellular protein implicated in nucleotide excision DNA repair. Withers-Ward E.S., Jowett J.B., Stewart S.A., Xie Y.M., Garfinkel A., Shibagaki Y., Chow S.A., Shah N., Hanaoka F., Sawitz D.G., Armstrong R.W., Souza L.M., Chen I.S.J. *J. Virol.* 71:9732-9742(1997) Two human homologs of Rad23 are functionally interchangeable in complex formation and stimulation of XPC repair activity. Sugasawa K., Ng J.M., Masutani C., Maekawa T., Uchida A., van der Spek P.J., Eker A.P., Rademakers S., Visser C., Aboussekhr A., Wood R.D., Hanaoka F., Bootsma D., Hoeijmakers J.H. *Mol. Cell. Biol.* 17:6924-6931(1997) Interaction of hHR23 with S5a. The ubiquitin-like domain of hHR23 mediates interaction with S5a subunit of 26 S proteasome. Hiyama H., Yokoi M., Masutani C., Sugasawa K., Maekawa T., Tanaka K., Hoeijmakers J.H., Hanaoka F. *J. Biol. Chem.* 274:28019-28025(1999) Ataxin-3, the MJD1 gene product, interacts with the two human homologs of yeast DNA repair protein RAD23, HHR23A and HHR23B. Wang G., Sawai N., Kotliarova S., Kanazawa I., Nukina N. *Hum. Mol. Genet.* 9:1795-1803(2000) Ubiquitin recognition by the DNA repair protein hHR23a. Wang Q., Goh A.M., Howley P.M., Walters K.J. *Biochemistry* 42:13529-13535(2003) Rad23 ubiquitin-associated domains (UBA) inhibit 26 S proteasome-catalyzed proteolysis by sequestering lysine 48-linked polyubiquitin chains. Raasi S., Pickart C.M. *J. Biol. Chem.* 278:8951-8959(2003) Binding of polyubiquitin chains to ubiquitin-associated (UBA) domains of HHR23A. Raasi S., Orlov I., Fleming K.G., Pickart C.M. *J. Mol. Biol.* 341:1367-1379(2004) Evidence for distinct functions for human DNA repair factors hHR23A and hHR23B. Chen L., Madura K. *FEBS Lett.* 580:3401-3408(2006) Ubiquitin receptor proteins hHR23a and hPLIC2 interact. Kang Y., Zhang N., Koepp D.M., Walters K.J. *J. Mol. Biol.* 365:1093-1101(2007) ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. Matsuoka S., Ballif B.A., Smogorzewska A., McDonald E.R. III, Hurov K.E., Luo J., Bakalarski C.E., Zhao Z., Solimini N., Lerenthal Y., Shiloh Y., Gygi S.P., Elledge S.J. *Science* 316:1160-1166(2007) Components of the ubiquitin-proteasome pathway compete for surfaces on Rad23 family proteins. Goh A.M., Walters K.J., Elsassser S., Verma R., Deshaies R.J., Finley D., Howley P.M. *BMC Biochem.* 9:4-4(2008) A quantitative atlas of mitotic phosphorylation. Dephoure N., Zhou C., Villen J., Beausoleil S.A., Bakalarski C.E., Elledge S.J., Gygi S.P. *Proc. Natl. Acad. Sci. U.S.A.* 105:10762-10767(2008) Quantitative phosphoproteomic analysis of T cell receptor signaling reveals system-wide modulation of protein-protein interactions. Mayya V., Lundgren D.H., Hwang S.-I., Rezaul K., Wu L., Eng J.K., Rodionov V., Han D.K. *Sci. Signal.* 2:RA46-RA46(2009) HIV-1 replication through hHR23A-mediated interaction of Vpr with 26S proteasome. Li G., Elder R.T., Dubrovsky L., Liang D., Pushkarsky T., Chiu K., Fan T., Sire J., Bukrinsky M., Zhao R.Y. *PLoS ONE* 5:E11371-E11371(2010) Quantitative phosphoproteomics reveals widespread full phosphorylation site occupancy during mitosis. Olsen J.V., Vermeulen M., Santamaria A., Kumar C., Miller M.L., Jensen L.J., Gnad F., Cox J., Jensen T.S., Nigg E.A., Brunak S., Mann M. *Sci. Signal.* 3:RA3-RA3(2010) Initial characterization of the human central proteome. Burkard T.R., Planyavsky M., Kaupe I., Breitwieser F.P., Buerckstuehmer T., Bennett K.L., Superti-Furga G., Colinge J. *BMC Syst. Biol.* 5:17-17(2011) System-wide temporal characterization of the proteome and phosphoproteome of human embryonic stem cell differentiation. Rigbolt K.T., Prokhorova T.A., Akimov V., Henningsen J., Johansen P.T., Kratchmarova I., Kassem M., Mann M., Olsen J.V., Blagoev B. *Sci. Signal.* 4:RS3-RS3(2011) An enzyme assisted RP-RPLC approach for in-depth analysis of human liver phosphoproteome. Bian Y., Song C., Cheng K., Dong M., Wang F., Huang J., Sun D., Wang L., Ye M., Zou H.J. *Proteomics* 96:253-262(2014) Structure of a human DNA repair protein UBA domain that interacts with HIV-1 Vpr. Dieckmann T., Withers-Ward E.S., Jarosinski M.A., Liu C.F., Chen I.S.Y., Feigon J. *Nat. Struct. Biol.* 5:1042-1047(1998) Biochemical and structural analysis of the interaction between the UBA(2) domain of the DNA repair protein HHR23A and HIV-1 Vpr. Withers-Ward E.S., Mueller T.D., Chen I.S.Y., Feigon J. *Biochemistry* 39:14103-14112(2000) Solution structures of UBA domains reveal a conserved hydrophobic surface for protein-protein interactions. Mueller T.D., Feigon J.J. *Mol. Biol.* 319:1243-1255(2002) Structural determinants for the binding of ubiquitin-like domains to the proteasome. Mueller T.D., Feigon J. *EMBO J.* 22:4634-4645(2003) DNA-repair protein hHR23a alters its protein structure upon binding proteasomal subunit S5a. Walters K.J., Lech P.J., Goh A.M., Wang Q., Howley P.M. *Proc. Natl. Acad. Sci. U.S.A.* 100:12694-12699(2003) Structure of the XPC binding domain of hHR23A reveals hydrophobic patches for protein interaction. Kamionka M., Feigon J. *Protein Sci.* 13:2370-2377(2004) Structural determinants for selective recognition of a Lys48-linked polyubiquitin chain by a UBA domain. Varadan R., Assfalg M., Raasi S., Pickart C., Fushman D. *Mol. Cell* 18:687-698(2005)

NCBI AND UNIPROT INFORMATION

NCBI GI #	392996949
NCBI GeneID	5886
NCBI Accession #	NP_001257291.1
NCBI GenBank Nucleotide #	NM_001270362.1
UniProt Accession #	P54725; Q59EU8; Q5M7Z1; K7ESE3
Molecular Weight	41.6 kDa
NCBI Official Full Name	UV excision repair protein RAD23 homolog A isoform 2
NCBI Official Synonym Full Names	RAD23 homolog A, nucleotide excision repair protein
NCBI Official Symbol	RAD23A
NCBI Official Synonym Symbols	HR23A; HHR23A
NCBI Protein Information	UV excision repair protein RAD23 homolog A
UniProt Protein Name	UV excision repair protein RAD23 homolog A
UniProt Gene Name	RAD23A
UniProt Synonym Gene Names	HR23A; hHR23A
UniProt Entry Name	RD23A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rad23a-recombinant-protein-e-coli-or-yeast-or-baculovirus-or-mammalian-cell/81657>