

Rat Anti-tissue transglutaminase IgA ELISA Kit | TTG-IgA elisa kit

Rat Anti-tissue transglutaminase IgA ELISA Kit

CATALOG #
AAA81950

REACTIVITY
Rat


PRODUCT SYNONYMS

Anti-tissue transglutaminase IgA; Rat Anti-tissue transglutaminase IgA ELISA Kit; TTG-IgA elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of TTG-IgA. No significant cross-reactivity or interference between TTG-IgA and analogues was observed. NOTE: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between TTG-IgA and all the analogues, therefore, cross reaction may still exist in some cases.
Samples	Serum, plasma, cell culture supernatants, body fluid and tissue homogenate
Assay Type	Quantitative Competitive
Sensitivity	1.0 ng/mL
Preparation and Storage	Store all reagents at 2-8 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TTG-IgA elisa kit	Intended Uses: This TTG-IgA ELISA kit is a 1.5 hour solid-phase ELISA designed for the quantitative determination of Rat TTG-IgA. This ELISA kit for research use only, not for therapeutic or test applications! Principle of the Assay: TTG-IgA ELISA kit applies the competitive enzyme immunoassay technique utilizing a polyclonal anti-TTG-IgA antibody and an TTG-IgA-HRP conjugate. The assay sample and buffer are incubated together with TTG-IgA-HRP conjugate in pre-coated plate for one hour. After the incubation period, the wells are decanted and washed five times. The wells are then incubated with a substrate for HRP enzyme. The product of the enzyme-substrate reaction forms a blue colored complex. Finally, a stop solution is added to stop the reaction, which will then turn the solution yellow. The intensity of color is measured spectrophotometrically at 450nm in a microplate reader. The intensity of the color is inversely proportional to the TTG-IgA concentration since TTG-IgA from samples and TTG-IgA-HRP conjugate compete for the anti-TTG-IgA antibody binding site. Since the number of sites is limited, as more sites are occupied by TTG-IgA from the sample, fewer sites are left to bind TTG-IgA-HRP conjugate. A standard curve is plotted relating the intensity of the color (O.D.) to the concentration of standards. The TTG-IgA concentration in each sample is interpolated from this standard curve.
Product Categories/Family for TTG-IgA elisa kit	Cell Biology

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/ttg-iga-rat-elisa-kits/anti-tissue-transglutaminase-iga/81950>