

Rat Phosphorylated Adenosine Monophosphate Activated Protein ELISA Kit | P-AMPAP elisa kit

Rat Phosphorylated Adenosine Monophosphate Activated Protein ELISA Kit

CATALOG # AAA82102	REACTIVITY Rat
------------------------------	--------------------------


PRODUCT SYNONYMS

Phosphorylated Adenosine Monophosphate Activated Protein; Rat Phosphorylated Adenosine Monophosphate Activated Protein ELISA Kit; P-AMPAP elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Samples	Serum, plasma, Cell Culture Supernatants, body fluid and tissue homogenate
Assay Type	Competitive
Sensitivity	1.0 pg/mL.
Intended Uses	This PPARGgamma ELISA kit is a 1.5 hour solid-phase ELISA designed for the quantitative determination of Rat PPARGgamma. This ELISA kit for research use only. not for therapeutic applications
Preparation and Storage	Store all reagents at 2-8 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for P-AMPAP elisa kit

Principle of the Assay: PPARGgamma ELISA kit applies the competitive enzyme immunoassay technique utilizing a monoclonal anti-PPARGgamma antibody and an PPARGgamma-HRP conjugate. The assay sample and buffer are incubated together with PPARGgamma-HRP conjugate in pre-coated plate for one hour. After the incubation period, the wells are decanted and washed five times. The wells are then incubated with a substrate for HRP enzyme. The product of the enzyme-substrate reaction forms a blue colored complex. Finally a stop solution is added to stop the reaction, which will then turn the solution yellow. The intensity of color is measured spectrophotometrically at 450nm in a microplate reader. The intensity of the color is inversely proportional to the PPARGgamma concentration since PPARGgamma from samples and PPARGgamma-HRP conjugate compete for the anti-PPARGgamma antibody binding site. Since the number of sites is limited, as more sites are occupied by PPARGgamma from the sample, fewer sites are left to bind PPARGgamma-HRP conjugate. A standard curve is plotted relating the intensity of the color (O.D.) to the concentration of standards. The PPARGgamma concentration in each sample is interpolated from this standard curve.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

