

Human Mannose Binding Lectin C ELISA Kit | MBL C elisa kit

Human Mannose Binding Lectin C ELISA Kit

CATALOG #
AAA85345

REACTIVITY
Human


PRODUCT SYNONYMS

Mannose Binding Lectin C; Human Mannose Binding Lectin C ELISA Kit; MBL C elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Samples	Serum, plasma, Cell Culture Supernatants, body fluid and tissue homogenate
Assay Type	Competitive
Detection Range	5.0-100ng/mL
Sensitivity	1.0ng/mL
Preparation and Storage	Store all reagents at 2-8 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MBL C elisa kit	Intended Uses: This MBL-C ELISA kit is intended Laboratory for research use only. The Stop Solution changes the color from blue to yellow and the intensity of the color is measured at 450 nm using a spectrophotometer. In order to measure the concentration of MBL-C in the sample, this MBL-CELISA Kit includes a set of calibration standards. The calibration standards are assayed at the same time as the samples and allow the operator to produce a standard curve of Optical Density versus MBL-C concentration. The concentration of MBL-C in the samples is then determined by comparing the O.D. of the samples to the standard curve. Principle of the Assay: This MBL-C enzyme linked immunosorbent assay applies a technique called a quantitative sandwich immunoassay. The microtiter plate provided in this kit has been pre-coated with a monoclonal antibody specific for MBL-C. Standards or samples are then added to the microtiter plate wells and MBL-C if present, will bind to the antibody pre-coated wells. In order to quantitatively determine the amount of MBL-C present in the sample, a standardized preparation of horseradish peroxidase (HRP)-conjugated polyclonal antibody, specific for MBL-C are added to each well to "sandwich" the MBL-C immobilized on the plate. The microtiter plate undergoes incubation, and then the wells are thoroughly washed to remove all unbound components. Next, A and B substrate solution is added to each well. The enzyme (HRP) and substrate are allowed to react over a short incubation period. Only those wells that contain MBL-C and enzyme-conjugated antibody will exhibit a change in colour. The enzyme-substrate reaction is terminated by the addition of a sulphuric acid solution and the colour change is measured spectrophotometrically at a wavelength of 450 nm.
Product Categories/Family for MBL C elisa kit	Immunology

NCBI AND UNIPROT INFORMATION

NCBI GI #	6754656
NCBI GeneID	17195
NCBI Accession #	NP_034906.1
NCBI GenBank Nucleotide #	NM_010776.1
Molecular Weight	25,957 Da
NCBI Official Full Name	mannose-binding protein C
NCBI Official Synonym Full Names	mannose-binding lectin (protein C) 2
NCBI Official Symbol	Mbl2
NCBI Official Synonym Symbols	MBL; L-MBP; MBL-C; MBP-C
NCBI Protein Information	mannose-binding protein C; RARF/P28A; mannan-binding protein; mannose binding lectin (C); RA-reactive factor P28A subunit; mannose binding lectin, liver (C)
UniProt Protein Name	Mannose-binding protein C
UniProt Gene Name	Mbl2
UniProt Synonym Gene Names	MBP-C; RARF/P28A
UniProt Entry Name	MBL2_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mbi-c-human-elisa-kits/mannose-binding-lectin-c/85345>