

Rabbit Cystatin C ELISA Kit | Cys-C elisa kit

Rabbit Cystatin C ELISA Kit

CATALOG #
AAA85472

REACTIVITY
Rabbit


PRODUCT SYNONYMS

Cystatin C; Rabbit Cystatin C ELISA Kit; Rabbit Cystatin C ELISA Kit (Cys-C); Cystatin C ELISA Kit (Rabbit); Cys-C elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rabbit
Assay Type	Sandwich
Preparation and Storage	Store all reagents at 2-8 degree C

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Cys-C elisa kit

Intended Uses: This CYS-CELISA kit is intended Laboratory for research use only. The Stop Solution changes the color from blue to yellow and the intensity of the color is measured at 450 nm using a spectrophotometer. In order to measure the concentration of CYS-C in the sample, this CYS-CELISA Kit includes a set of calibration standards. The calibration standards are assayed at the same time as the samples and allow the operator to produce a standard curve of Optical Density versus CYS-C concentration. The concentration of CYS-C in the samples is then determined by comparing the O.D. of the samples to the standard curve.

Principle of the Assay: This CYS-C enzyme linked immunosorbent assay applies a technique called a quantitative sandwich immunoassay. The microtiter plate provided in this kit has been pre-coated with a monoclonal antibody specific for CPP.HS. Standards or samples are then added to the microtiter plate wells and CYS-C if present, will bind to the antibody pre-coated wells. In order to quantitatively determine the amount of CYS-C present in the sample, a standardized preparation of horseradish peroxidase (HRP)-conjugated polyclonal antibody, specific for CYS-C are added to each well to "sandwich" the CYS-C immobilized on the plate. The microtiter plate undergoes incubation, and then the wells are thoroughly washed to remove all unbound components. Next, A and B substrate solution is added to each well. The enzyme (HRP) and substrate are allowed to react over a short incubation period. Only those wells that contain CYS-C and enzyme-conjugated antibody will exhibit a change in colour. The enzyme-substrate reaction is terminated by the addition of a sulphuric acid solution and the colour change is measured spectrophotometrically at a wavelength of 450 nm.

NCBI AND UNIPROT INFORMATION

NCBI GI #	296643
NCBI GeneID	1471
UniProt Accession #	P01034; Q6FGW9; B2R5J9; D3DW42
Molecular Weight	15,799 Da
NCBI Official Full Name	cystatin C
NCBI Official Synonym Full Names	cystatin C
NCBI Official Symbol	CST3
NCBI Official Synonym Symbols	ARMD11
NCBI Protein Information	cystatin-C; cystatin 3; cystatin-3; gamma-trace; post-gamma-globulin; bA218C14.4 (cystatin C); neuroendocrine basic polypeptide
UniProt Protein Name	Cystatin-C
UniProt Gene Name	CST3
UniProt Entry Name	CYTC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.