

Mouse Macrophage chemotactic factor ELISA Kit | MCF elisa kit

Mouse Macrophage chemotactic factor ELISA Kit

CATALOG #
AAA87481

REACTIVITY
Mouse


PRODUCT SYNONYMS

Macrophage chemotactic factor; Mouse Macrophage chemotactic factor ELISA Kit; MCF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Assay Type	Sandwich
Preparation and Storage	Store all reagents at 2-8 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MCF elisa kit

Intended Uses: This MDC ELISA kit is intended Laboratory for research use only. The Stop Solution changes the color from blue to yellow and the intensity of the color is measured at 450 nm using a spectrophotometer. In order to measure the concentration of MDC in the sample, this MDC ELISA Kit includes a set of calibration standards. The calibration standards are assayed at the same time as the samples and allow the operator to produce a standard curve of Optical Density versus MDC concentration. The concentration of MDC in the samples is then determined by comparing the O.D. of the samples to the standard curve.

Principle of the Assay: This MDC enzyme linked immunosorbent assay applies a technique called a quantitative sandwich immunoassay. The microtiter plate provided in this kit has been pre-coated with a monoclonal antibody specific for MDC. Standards or samples are then added to the microtiter plate wells and MDC if present, will bind to the antibody pre-coated wells. In order to quantitatively determine the amount of MDC present in the sample, a standardized preparation of horseradish peroxidase (HRP)-conjugated polyclonal antibody, specific for MDC are added to each well to "sandwich" the MDC immobilized on the plate. The microtiter plate undergoes incubation, and then the wells are thoroughly washed to remove all unbound components. Next, A and B substrate solution is added to each well. The enzyme (HRP) and substrate are allowed to react over a short incubation period. Only those wells that contain MDC and enzyme-conjugated antibody will exhibit a change in colour. The enzyme-substrate reaction is terminated by the addition of a sulphuric acid solution and the colour change is measured spectrophotometrically at a wavelength of 450 nm.

NCBI AND UNIPROT INFORMATION

NCBI GI #	484255
Molecular Weight	57,911 Da
NCBI Official Full Name	MDC protein
UniProt Protein Name	Disintegrin and metalloproteinase domain-containing protein 11
UniProt Gene Name	ADAM11
UniProt Synonym Gene Names	MDC; ADAM 11; MDC
UniProt Entry Name	ADA11_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mcf-mouse-elisa-kits/macrophage-chemotactic-factor/87481>