

Porcine Activated protein C ELISA Kit | APC elisa kit

Porcine Activated protein C ELISA Kit

CATALOG #
AAA88002

REACTIVITY
Porcine


PRODUCT SYNONYMS

Activated protein C; Porcine Activated protein C ELISA Kit; APC elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Porcine
Specificity	This assay has high sensitivity and excellent specificity for detection of APC. No significant cross-reactivity or interference between APC and analogues was observed. NOTE: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between APC and all the analogues, therefore, cross reaction may still exist in some cases.
Samples	Serum, plasma, cell culture supernatants, body fluid and tissue homogenate
Assay Type	Quantitative Sandwich
Sensitivity	0.1 ng/mL
Preparation and Storage	Store all reagents at 2-8 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for APC elisa kit	Intended Uses: This APC ELISA kit is a 1.5 hour solid-phase ELISA designed for the quantitative determination of Porcine APC. This ELISA kit for research use only, not for therapeutic or test applications! Principle of the Assay: APC ELISA kit applies the quantitative sandwich enzyme immunoassay technique. The microtiter plate has been pre-coated with a monoclonal antibody specific for APC. Standards or samples are then added to the microtiter plate wells and APC if present, will bind to the antibody pre-coated wells. In order to quantitatively determine the amount of APC present in the sample, a standardized preparation of horseradish peroxidase (HRP)-conjugated polyclonal antibody, specific for APC are added to each well to "sandwich" the APC immobilized on the plate. The microtiter plate undergoes incubation, and then the wells are thoroughly washed to remove all unbound components. Next, substrate solutions are added to each well. The enzyme (HRP) and substrate are allowed to react over a short incubation period. Only those wells that contain APC and enzyme-conjugated antibody will exhibit a change in color. The enzyme-substrate reaction is terminated by addition of a sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450 nm. A standard curve is plotted relating the intensity of the color (O.D.) to the concentration of standards. The APC concentration in each sample is interpolated from this standard curve.
Product Categories/Family for APC elisa kit	Cardiovascular

NCBI AND UNIPROT INFORMATION

NCBI GI #	53759122
NCBI GeneID	324
NCBI Accession #	NP_000029.2
NCBI GenBank Nucleotide #	NM_000038.5
Molecular Weight	300,439 Da
NCBI Official Full Name	adenomatous polyposis coli protein isoform b
NCBI Official Synonym Full Names	adenomatous polyposis coli
NCBI Official Symbol	APC
NCBI Official Synonym Symbols	GS; DP2; DP3; BTPS2; DP2.5; PPP1R46
NCBI Protein Information	adenomatous polyposis coli protein; deleted in polyposis 2.5; adenomatosis polyposis coli tumor suppressor; protein phosphatase 1, regulatory subunit 46
UniProt Protein Name	Adenomatous polyposis coli protein
UniProt Gene Name	APC
UniProt Synonym Gene Names	DP2.5; Protein APC
UniProt Entry Name	APC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.