

Chicken Low density lipoprotein immune complex ELISA Kit | LDL-IC elisa kit

Chicken Low density lipoprotein immune complex ELISA Kit

CATALOG #
AAA96751

REACTIVITY
Chicken


PRODUCT SYNONYMS

Low density lipoprotein immune complex; Chicken Low density lipoprotein immune complex ELISA Kit; LDL-IC elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Chicken
Samples	Serum, plasma, Cell Culture Supernatants, body fluid and tissue homogenate
Assay Type	Competitive
Sensitivity	0.1 ng/mL.
Preparation and Storage	Store all reagents at 2-8 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for LDL-IC elisa kit

Principle of the assay: LDL-IC ELISA kit applies the competitive enzyme immunoassay technique utilizing a monoclonal anti-LDL-IC antibody and an LDL-IC-HRP conjugate. The assay sample and buffer are incubated together with LDL-IC-HRP conjugate in pre-coated plate for one hour. After the incubation period, the wells are decanted and washed five times. The wells are then incubated with a substrate for HRP enzyme. The product of the enzyme-substrate reaction forms a blue colored complex. Finally, a stop solution is added to stop the reaction, which will then turn the solution yellow. The intensity of color is measured spectrophotometrically at 450nm in a microplate reader. The intensity of the color is inversely proportional to the LDL-IC concentration since LDL-IC from samples and LDL-IC-HRP conjugate compete for the anti-LDL-IC antibody binding site. Since the number of sites is limited, as more sites are occupied by LDL-IC from the sample, fewer sites are left to bind LDL-IC-HRP conjugate. A standard curve is plotted relating the intensity of the color (O.D.) to the concentration of standards. The LDL-IC concentration in each sample is interpolated from this standard curve.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.