

Rat Neutrophil Activating Protein-2 ELISA Kit | NAP-2 elisa kit

Rat Neutrophil Activating Protein-2 ELISA Kit

CATALOG #
AAA97928

REACTIVITY
Rat


PRODUCT SYNONYMS

Neutrophil Activating Protein-2; Rat Neutrophil Activating Protein-2 ELISA Kit; NAP-2/CXCL7/CTAP III/PPBP/connective tissue-activating peptide III/CTAP3CTAP-III/CXC chemokine ligand 7/C-X-C motif chemokine 7/CXCL7b-TG1/LA-PF4/LDGFTC1/Leukocyte-derived growth factor/low-affinity platelet factor IV/Macrophage-derived growth factor/MDGFTC2/NAP-2-L1/PBPTHGB/platelet basic protein/PPBP/pro-platelet basic protein (chemokine (C-X-C motif) ligand 7)/SCAR10/SCYB7beta-thromboglobulin/small inducible cytokine B7/small inducible cytokine subfamily B; member 7/Small-inducible cytokine B7/TGB/TGB1B-TG1/THBGB1/thrombocidin 1/thrombocidin 2/thromboglobulin; beta-1; NAP-2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of NAP-2. No significant cross-reactivity or interference between NAP-2 and analogues was observed.
Sequence Length	70
Assay Type	Sandwich
Samples	Serum, plasma, tissue homogenates and other biological fluids
Detection Range	0.156-10ng/ml
Sensitivity	0.094ng/ml
Intra-assay Precision	CV<8%
Inter-assay Precision	CV<10%
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NAP-2 elisa kit	Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Capture antibody was pre-coated onto 96-well plates. And the biotin conjugated antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and washed with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the target amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of target can be calculated.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	284434905
NCBI GeneID	5473
Molecular Weight	13,894 Da
NCBI Official Full Name	neutrophil-activating protein 2, partial
NCBI Official Synonym Full Names	pro-platelet basic protein
NCBI Official Symbol	PPBP
NCBI Official Synonym Symbols	PBP; TC1; TC2; TGB; LDGF; MDGF; TGB1; B-TG1; CTAP3; CXCL7; NAP-2; SCYB7; THBGB; LA-PF4; THBGB1; Beta-TG; CTAPIII; CTAP-III
NCBI Protein Information	platelet basic protein
UniProt Protein Name	Platelet basic protein
UniProt Gene Name	PPBP
UniProt Synonym Gene Names	CTAP3; CXCL7; SCYB7; TGB1; THBGB1; PBP; LDGF; MDGF; CTAP-III; CTAP-III(1-81); Beta-TG; NAP-2(74); NAP-2(73); NAP-2; NAP-2(1-66); NAP-2(1-63)
UniProt Entry Name	CXCL7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/nap-2-rat-elisa-kits/neutrophil-activating-protein-2/97928>