

Mouse Soluble Interleukin-2 Receptor Alpha Chain ELISA Kit | IL-2sRalpha/CD25 elisa kit

Mouse Soluble Interleukin-2 Receptor Alpha Chain ELISA Kit

CATALOG #
AAA98526

REACTIVITY
Mouse


PRODUCT SYNONYMS

Soluble Interleukin-2 Receptor Alpha Chain; Mouse Soluble Interleukin-2 Receptor Alpha Chain ELISA Kit; IL-2sRalpha/CD25/IL-2 R alpha/IL2RA/CD25/CD25 antigen/IDDM10/IL-2 receptor subunit alpha/IL2R/IL-2R subunit alpha/IL-2-RA/IL2-RA/interleukin 2 receptor; alpha/interleukin-2 receptor subunit alpha/p55/TAC antigen/TCGFR ; IL-2sRalpha/CD25 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of IL-2sR α /CD25 . No significant cross-reactivity or interference between IL-2sR α /CD25 and analogues was observed. Note: Limited by current skills and knowledge, it is difficult for us to complete the crossreactivity detection between IL-2sR α /CD25 and all the analogues, therefore, cross reaction may still exist.
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	78.125-5000pg/ml
Sensitivity	46.875pg/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level IL- 2sR α /CD25 were tested 20 times on one plate, respectively. Intra-Assay: CV<8%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level IL- 2sR α /CD25 were tested on 3 different plates, 8 replicates in each plate. CV (%) = SD/meanX100. Inter-Assay: CV<10%
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL-2sRalpha/CD25 elisa kit

Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti- IL-2sR α /CD25 antibody was pre-coated onto 96-well plates. And the biotin conjugated anti- IL-2sR α /CD25 antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and washed with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the IL-2sR α /CD25 amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of IL- 2sR α /CD25 can be calculated.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/il-2sralpha-cd25-mouse-elisa-kits/soluble-interleukin-2-receptor-alpha-chain/98526>