

Human T-lymphocyte activation antigen CD86 ELISA Kit | CD86 elisa kit

Human T-lymphocyte activation antigen CD86 ELISA Kit

CATALOG #
AAA98775

REACTIVITY
Human


PRODUCT SYNONYMS

T-lymphocyte activation antigen CD86; Human T-lymphocyte activation antigen CD86 ELISA Kit; B7-2/CD86/Activation B7-2 antigen/B70/B7-2 antigen/B-lymphocyte activation antigen B7-2/BU63/CD28 antigen ligand 2/CD28LG2/CD86 antigen/CD86 molecule/CTLA-4 counter-receptor B7.2/FUN-1/LAB72/T-lymphocyte activation antigen CD86; CD86 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	247
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	0.156-10ng/ml
Sensitivity	0.094ng/ml
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES



Standard Curve (Sample)

Results of a typical standard run of a Human CD86 ELISA Kit are shown below. This standard curve was generated at our lab for demonstration purpose only. Each user should obtain their own standard curve as per experiment. (N/A=not applicable)

X	ng/ml	0	0.156	0.313	0.625	1.25	2.5	5	10
Y	OD450								

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CD86 elisa kit

Principle of the Assay||This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-Human CD86 antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-Human CD86 antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and wash with wash buffer. HRPStreptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the Human CD86 amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of Human CD86 can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	332634954
NCBI GeneID	942
NCBI Accession #	NP_001193854.1
NCBI GenBank Nucleotide #	NM_001206925.1
Molecular Weight	28,420 Da
NCBI Official Full Name	T-lymphocyte activation antigen CD86 isoform 5
NCBI Official Synonym Full Names	CD86 molecule
NCBI Official Symbol	CD86
NCBI Official Synonym Symbols	B70; B7-2; B7.2; LAB72; CD28LG2
NCBI Protein Information	T-lymphocyte activation antigen CD86
UniProt Protein Name	T-lymphocyte activation antigen CD86
UniProt Gene Name	CD86
UniProt Synonym Gene Names	CD28LG2
UniProt Entry Name	CD86_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/t-lymphocyte-activation-antigen-cd86/98775>