

Rat Sclerostin ELISA Kit | Sost elisa kit

Rat Sclerostin ELISA Kit

CATALOG #
AAA99012

REACTIVITY
Rat

PRODUCT SYNONYMS

Sclerostin; Rat Sclerostin ELISA Kit; Sost/SOST (Sclerostin); Sost elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Sost. No significant cross-reactivity or interference between Sost and analogues was observed.
Sequence Length	213
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	31.25-2000pg/ml
Sensitivity	18.75pg/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Sost were tested 20 times on one plate, respectively. Intra-Assay: CV<8%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Sost were tested on 3 different plates, 8 replicates in each plate. CV (%) = SD/meanX100. Inter-Assay: CV<10%
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES

Typical Data & Standard Curve

Results of a typical standard run of a Sost ELISA Kit are shown below. This standard curve was generated at our lab for demonstration purpose only. Each user should obtain their own standard curve as per experiment. (N/A=not applicable)

X	pg/ml	0	31.25	62.5	125	250	500	1000	2000
Y	OD450	0.068	0.269	0.407	0.648	1.035	1.453	1.907	2.589

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Sost elisa kit

Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti- Sost antibody was pre-coated onto 96-well plates. And the biotin conjugated anti- Sost antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and wash with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the Sost amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of Sost can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	13376846
NCBI GeneID	50964
NCBI Accession #	NP_079513.1
NCBI GenBank Nucleotide #	NM_025237.2
Molecular Weight	24,264 Da
NCBI Official Full Name	sclerostin
NCBI Official Synonym Full Names	sclerostin
NCBI Official Symbol	SOST
NCBI Official Synonym Symbols	CDD; VBCH; SOST1
NCBI Protein Information	sclerostin
UniProt Protein Name	Sclerostin
UniProt Gene Name	SOST
UniProt Entry Name	SOST_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/sost-rat-elisa-kits/sclerostin/99012>