

Rat Coagulation Factor VII ELISA Kit | F7 elisa kit

Rat Coagulation Factor VII ELISA Kit

CATALOG #
AAA99111

REACTIVITY
Rat

PRODUCT SYNONYMS

Coagulation Factor VII; Rat Coagulation Factor VII ELISA Kit; F7 (Coagulation Factor VII)/FVII/Proconvertin/SPCA; F7 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of F7. No significant crossreactivity or interference between F7 and analogues was observed.
Sequence Length	21
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	62.5-4000pg/ml
Sensitivity	37.5pg/ml
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level F7 were tested 20 times on one plate, respectively. Intra-Assay: CV<8%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level F7 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<10%
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES

Typical Data & Standard Curve									
Results of a typical standard run of a F7 ELISA Kit are shown below. This standard curve was generated at our lab for demonstration purpose only. Each user should obtain their own standard curve as per experiment. (N/A: not applicable)									
X	pg/ml	0	62.5	125	250	500	1000	2000	4000
Y	OD450	0.01	0.072	0.116	0.268	0.585	1.135	1.883	2.313

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for F7 elisa kit

Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti- F7 antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-F7 antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and wash with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the F7 amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of F7 can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	12750767
NCBI GeneID	2155
Molecular Weight	49,320 Da
NCBI Official Full Name	coagulation factor VII, partial
NCBI Official Synonym Full Names	coagulation factor VII
NCBI Official Symbol	F7
NCBI Official Synonym Symbols	SPCA
NCBI Protein Information	coagulation factor VII
UniProt Protein Name	Coagulation factor VII
UniProt Gene Name	F7
UniProt Synonym Gene Names	SPCA
UniProt Entry Name	FA7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/f7-rat-elisa-kits/coagulation-factor-vii/99111>