

Human T-cell surface antigen CD2 ELISA Kit | CD2 elisa kit

Human T-cell surface antigen CD2 ELISA Kit

CATALOG #
AAA99626

REACTIVITY
Human


PRODUCT SYNONYMS

T-cell surface antigen CD2; Human T-cell surface antigen CD2 ELISA Kit; LFA-2/CD2 (T-cell surface antigen CD2)/T11/CD2 antigen/CD2 antigen (p50); sheep red blood cell receptor/CD2 molecule/Erythrocyte receptor/FLJ46032/LFA-2/LFA-3 receptor/lymphocyte-function antigen-2/Rosette receptor/SRBC/T11/T-cell surface antigen CD2; CD2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	351
Assay Type	Sandwich ELISA, Double Antibody
Samples	Serum, Plasma, Cell Culture Supernatant, cell or tissue lysate, Other liquid samples
Detection Range	0.313-20ng/ml
Sensitivity	0.188ng/ml
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES



Standard Curve (Sample)

Typical Data & Standard Curve

Results of a typical standard run of a Human LFA-2 ELISA Kit are shown below. This standard curve was generated at our lab for demonstration purpose only. Each user should obtain their own standard curve as per experiment. (N/A=not applicable)

X	ng/ml	0	0.313	0.625	1.25	2.5	5	10	20
Y	OD450	0.082	0.171	0.254	0.409	0.704	1.091	1.587	2.025

Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CD2 elisa kit

Principle of the Assay||This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-Human LFA-2 antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-Human LFA-2 antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and wash with wash buffer. HRP-Streptavidin was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the Human LFA-2 amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of Human LFA-2 can be calculated.

NCBI AND UNIPROT INFORMATION

NCBI GI #	156071472
NCBI GeneID	914
NCBI Accession #	NP_001758.2
NCBI GenBank Nucleotide #	NM_001767.3
Molecular Weight	39,448 Da
NCBI Official Full Name	T-cell surface antigen CD2
NCBI Official Synonym Full Names	CD2 molecule
NCBI Official Symbol	CD2
NCBI Official Synonym Symbols	T11; SRBC; LFA-2
NCBI Protein Information	T-cell surface antigen CD2
UniProt Protein Name	T-cell surface antigen CD2
UniProt Gene Name	CD2
UniProt Synonym Gene Names	SRBC
UniProt Entry Name	CD2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/t-cell-surface-antigen-cd2/99626>